

**NYSDOH-NYSED Approved Opioid Overdose Prevention Training
Post-Test and Training Skills Checklist for Libraries**

Trainee Name _____ **Date** _____

Directions: Complete this Post-Test. If you have not achieved 100% accuracy, review the training and take the test again. Take your completed Post-Test/Training Checklist to the trainer representing the NYSDOH Registered Opioid Overdose Prevention Program who will review how to administer the naloxone products used in your facility and sign your certificate of completion. It is valid for 2 years.

The “Community Access Naloxone Law” allows trained non-medical persons to administer naloxone to another person to prevent an opioid/heroin overdose from becoming fatal.

T or F

Libraries may permit volunteer employees to be trained by an individual representing a Registered Opioid Overdose Prevention Program to administer an opioid antagonist in the event of an emergency under Public Health Law section 3309.

T or F

Which NYS agency laws, regulations or policies must be followed to provide and Opioid Overdose Prevention Program in schools or school district? (Choose all that apply)

- A. Public Health Laws/Regulations
- B. B. Education Laws/Regulations
- C. C. Public library policy/procedures

In an opioid overdose, the individual becomes sedated and loses the urge to breathe.

T or F

Naloxone is an opioid antagonist that reverses the signs of overdose and restores breathing. It works in:

- A. 2- 3 minutes
- B. 8-15 minutes
- C. 30-60 minutes

Signs of an opioid overdose include which of the following? (Choose all that apply)

- A. Unresponsive, body limp with slow or no breathing
- B. Lips and fingernails turn blu(ish)
- C. Agitation and Hyperactivity

Fentanyl is a much more powerful opioid than heroin creating a greater risk for overdose.

T or F

Which steps below are part of emergency treatment of a suspected opioid overdose? (Choose All that apply)

- A. Call 911, state the person is not breathing and activate School Emergency Response
- B. Determine if the person is conscious or unresponsive-shake, shout, perform sternal rub
- C. Follow emergency response (CPR/Rescue Breaths) and if suspected overdose GIVE NALOXONE
- D. Roll the person into recovery position and monitor breathing until EMS arrives
- E. Report administration of naloxone to Program/Clinical Director

NYSDOH-NYSED Approved Opioid Overdose Prevention Training Skills Checklist

Opioid Overdose Prevention Training Skills Checklist for Schools

Directions to trainer: Review the items with the trainee. Upon successful completion add trainees name to the Opioid Overdose Prevention Staff Training Log Summary and sign their certificate of completion.

Trainee Name:	
Trainer Name:	Date:
Trainer Title : <input type="checkbox"/> RN <input type="checkbox"/> NP <input type="checkbox"/> PA <input type="checkbox"/> MD/DO	
Library Name:	
NYSDOH Registered Opioid Overdose Program Name: <input type="checkbox"/> NA	

Knowledge Check Completed By Trainee	Trainer Name	Date
Provides a copy of Opioid Overdose Prevention Training post-test which indicates 100% accuracy		
Identifies location and contents of naloxone kit		
Describes documentation and reporting process per policy		

Skills Check for Administering Naloxone	Trainer Name	Date
Assesses the victim by Shake, Shout, Sternal Rub		
Calls/instructs someone to call 911 and activate school emergency response per policy		
Obtains and correctly administers naloxone following manufacturer's directions or steps below for Narcan® Nasal Spray		
<input type="checkbox"/> Take Narcan® Nasal Spray out of box. Peel back tab with the circle to open. <input type="checkbox"/> Hold the device with your thumb on the bottom of the plunger and 2 fingers on the nozzle. Do NOT press the plunger. <input type="checkbox"/> Tilt the person's head back and provide support under the neck with your hand. <input type="checkbox"/> insert the tip of the nozzle into one nostril, until your fingers on either side of the nozzle are against the bottom of the person's nose <input type="checkbox"/> Press the plunger firmly to give the dose of Narcan® Nasal Spray. <input type="checkbox"/> Remove the Narcan® Nasal Spray from nostril after giving the dose. <input type="checkbox"/> Place victim in recovery position. <input type="checkbox"/> If no or minimal response after 2-3 minutes, repeat with other dose of Narcan® Nasal Spray in box. <input type="checkbox"/> If not breathing, provide two slow breaths, then one rescue breath every 5 seconds.		

**Training is valid for 2 years. Annual review is highly recommended.
Maintain this record for 2 years unless superseded by updated training.**